

Clinical Order Set

Demographics

LTC Hypodermoclysis (Module)

(Subcutaneous Fluid Infus	ion)				
	Page 1 of 1			Kov	
Key: Req – Requisition MAR – Media	cation Administration Record K	– Kardex Dis – Dis	continued	Key	Phase
Instructions for completing this order set: ☑ Indicates a pre-selected order. To delete a pre-selected order, draw a line through it ☐ Must tick the box for order to be implemented. Orders not checked will not be implemented _ Fill in blank spaces as needed/appropriate - Indicates an item for consideration by Provider; is NOT an order LTC Hypodermoclysis (Module)					Module)
LTC Hypodermocrysis (ivid	<u>luulej</u>) s
Inclusion criteria: - Unable to ingest sufficient amounts of great service and the control of t	n for a short trial of rehydration, wit pulmonary edema		ime frame (48 to 72 hours)		LTC Hypodermoclysis (Module)
Communication Orders ✓ Refer to Protocol/Guideline/Standard Hypodermoclysis) for Long Term Care"	l, Nurse to follow <u>"10.3.16PR Pare</u>	nteral Infusion Ther	rapy (Subcutaneous Infusion	=	ГС НУБ
Patient Care					
Subcutaneous Butterfly Insertion, Mo	onitoring and site selection as per p	procedure			
Continuous Infusions Ordering considerations: - Maximum amount of fluid PER SITE is 2 - Two infusion sites are required for volu - Solution CANNOT be mixed with any m	me greater than 2,000 mL		ent		
IV SOLUTION					
☐ sodium chloride 0.9%☐ Ringers Lactate					
DURATION ☐ 24 hours ☐ 48 hours ☐ 72 hours					
VOLUME / ROUTE / DIRECTIONS					
\square 1,000 mL, SUBCUT, Titrate, Infuse 1,0	00 mL total volume per 24 hours				
\square 1,500 mL, SUBCUT, Titrate, Infuse 1,5	00 mL total volume per 24 hours				
\square 2,000 mL, SUBCUT, Titrate, Infuse 2,0	00 mL total volume per 24 hours				
☑ Goal is to infuse IV fluids via gravity p	er procedure; The body will preve	nt over-infusion int	o the third space		1
Signature, Designation	College License #	Date	Time Pag	e 1/1	