



Does the wound bed need a non-adherent contact layer?

- Is the wound bed fragile and requires protection?
- Is the wound painful?
- Are the dressing changes causing trauma to the wound bed?
- Is the exudate minimal and there is a concern that the cover (secondary) dressing will stick?



UrgoTul Ag Silver

- Anti-inflammatory
- Indicated for Acute and Chronic Wounds
- Must be in contact with the wound bed
- First application—1 to 3 days, then can be left in place up to 7 days or as indicated by treatment plan.



Inadine

- Pro-Inflammatory
- Indicated for Chronic, Slow healing and/or maintenance wounds.
- Must be in contact with the wound bed.
- Colour fading of product will indicate wear time



Bactigras

- Neutral (no inflammatory response)
- Chlorhexidine based
- Wear time is based on exudate level, usually between 2-4 days.
- Must be in contact with the wound bed



**Use all products in a single layer
Do not mix products**



Jelonet

- Paraffin based
- Do not over lap the periwound skin
- Daily dressing changes (it dries out)



Adaptic

- Petroleum based
- Overlap 2 cm of periwound
- Can stay in place up to 7 days
- Change as indicated by treatment plan



Mepitel

- Silicone based
- Cover 2 cm of periwound skin
- Can be left in place for 14 days
- Change as indicated by treatment plan



For detailed product information go to www.clwk.ca